

Table III - Intra-rater reliability of the longus colli muscle cross-sectional area in women with and without migraine.

Groups	Measure	Rest		Contraction			
		ICC	Limits of agreement (CI 95%)		ICC		
			Lower	Upper			
Migraine	CSA R	0.78	-0.19 (-0.294 to -0.086)	0.175 (0.071 to 0.279)	0.79	-0.182 (-0.284 to -0.080) 0.177 (0.075 to 0.279)	
	CSA L	0.68	-0.173 (-0.282 to -0.065)	0.207 (0.096 to 0.315)	0.53	-0.223 (-0.344 to -0.102) 0.201 (0.081 to 0.322)	
Control	CSA R	0.83	-0.177 (-0.291 to -0.063)	0.119 (0.006 to 0.233)	0.83	-0.138 (-0.216 to -0.061) 0.063 (-0.013 to 0.141)	
	CSA L	0.93	-0.093 (-0.147 to -0.039)	0.048 (-0.005 to 0.102)	0.35	-0.368 (-0.698 to -0.039) 0.493 (0.164 to 0.823)	

ICC (Intraclass correlation); CI 95% (confidence interval 95%); CSA (cross-sectional area - cm²); R (right); L (left).

Table IV - Inter-rater reliability of the longus colli muscle cross-sectional area in women with and without migraine.

Groups	Measures	Rest		Contraction			
		ICC	Limits of agreement (CI 95%)		ICC		
			Lower	Upper			
Migraine	CSA R	0.78	-0.282 (-0.411 to -0.152)	0.173 (0.044 to 0.303)	0,81	-0.206 (-0.329 to -0.083) 0.224 (0.102 to 0.347)	
	CSA L	0.78	-0.239 (-0,328 to -0.137)	0.104 (0.008 to 0.200)	0,89	-0.1(-0.172 to -0.028) 0.152 (0.080 to 0.224)	
Control	CSA R	0.86	-0.141 (-0.247 to -0.035)	0.136 (0.030 to 0.242)	0,90	-0.043 (-0.106 to 0.020) 0.123 (0.059 to 0.186)	
	CSA L	0.77	-0.203 (-0.319 to -0.043)	0.133 (0.166 to 0.442)	0,71	-0.129 (-0.264 to 0.005) 0.224 (0.889 to 0.359)	

ICC (Intraclass correlation); CI 95% (confidence interval 95%); CSA (cross-sectional area - cm²); R (right); L (left).